

Variety sheet

Lavanttaler Bananenapfel

Origin

Unknown cross



Synonyms

American Mother

Agronomic Profile

Flowering: medium, diploid

Vigour: medium

Susceptibility: n. a.

Productivity: Productivity medium, tends to biennial bearing, no preharvest fruit drop

Harvest time: 6-4 Week before Golden Delicious

Fruit

Size and shape: medium, globular-conoid

Ground and overcolour: Ground colour green; Overcolour dark red, compact, washed out, mottled; high (75%)

Susceptibility to russetting: low rustiness in the calyx/eye basin and/or in the stalk cavity (<5%)

Firmness of the pulp: 7,4-8,3 kg/cm² at harvest

Tendency to browning of pulp: n. a.

Sugar content: 12,0-13,9 °Brix at harvest

Acid content: 5,5-6,6 g/L MA at harvest

Specific weight: 0,82

Total content of Polyphenols¹: n. a.

TAC¹: n. a.

Content of Anthocyanin²: n. a.

1 - after two months of normal cool storage
2 - only for red fleshed varieties

12.04.2026 - www.laimburg.it

Vitamin C¹: n. a.

Pectin¹: n. a.

Flavour: n. a.

Storage: n. a.

Characteristics: n. a.

Juice production

Yield: n. a.

Stability of the colour: n. a.

Sugar content: n. a.

Acid content: n. a.

pH: n. a.

Total content of Polyphenols: n. a.

TAC: n. a.

Content of Anthocyanin²: n. a.

Vitamin C: n. a.

1 - after two months of normal cool storage
2 - only for red fleshed varieties

12.04.2026 - www.laimburg.it

